

THE CANADIAN ENTOMOLOGIST

Index to Volume 129 (1997)



THE ENTOMOLOGICAL SOCIETY OF CANADA, OTTAWA

- Abou-Zaid, M.M.** See Nicol et al., 991.
- Adler, P.H., and P.G. Mason.** Black flies (Diptera: Simuliidae) of east-central Saskatchewan, with description of a new species and implications for pest management, 81.
- Alarie, Y.** Taxonomic revision and phylogenetic analysis of the genus *Oreodytes* Seidlitz (Coleoptera: Dytiscidae: Hydroporinae) based on larval morphology, 399.
- Alford, A.R.** See Silk et al., 1001.
- Alhumeyri, A.** See AliNiazee et al., 27.
- AliNiazee, M.T., A. Alhumeyri, and M. Saeed.** Laboratory and field evaluation of a neem insecticide against *Archips rosanus* L. (Lepidoptera: Tortricidae), 27.
- Arnason, J.T.** See Nicol et al., 991.
- Asis, J.D.** See Torres et al., 1067.
- Bakke, A.** See Hui and Bakke, 579.
- Barclay, H.J.** Assessing natural selection in white pine weevils (*Pissodes strobi* Peck) (Coleoptera: Curculionidae) for overcoming resistance in trees: An evolutionary model, 1105.
- Barszcz, E.S.** See Stevenson and Barszcz, 187.
- Bauce, É.** See Trudel et al., 197.
- Behan-Pelletier, V.M.** The semiaquatic genus *Tegeocranellus* (Acari: Oribatida: Ameronothroidea) of North and Central America, 537.
- Bélanger, G.** See Pelletier et Bélanger, 801.
- Bellows, T.S., Jr.** See Headrick et al., 335.
- Blodgett, S.L., P.M. Denke, M.A. Ivie, C.W. O'Brien, and A.W. Lenssen.** *Listronotus montanus* Dietz (Coleoptera: Curculionidae) damaging spring wheat in Montana, 377.
- Bodnaryk, R.P.** See Palaniswamy et al., 43.
- Boiteau, G., and W.P.L. Osborn.** Behavioural effects of imidacloprid, a new nicotiny insecticide, on the potato aphid, *Macrosiphum euphorbiae* (Thomas) (Homoptera, Aphididae), 241.
- Boivin, G.** See Huber et al., 959.
- Bonduriensky, R., and R.J. Brooks.** A technique for measuring and marking live flies, 827.
- Borden, J.H.** See Partridge and Borden, 899.
- Bouchard, A.** See Brodeur et al., 1.
- Bousquet, Y., and S. Laplante.** Taxonomic review of the New World Pogonini (Coleoptera: Carabidae), 699. (See Errata.)
- Bown, T.A.** See Mitchell et al., 745.
- Broadbent, A.B., and D.J. Pree.** Resistance to insecticides in populations of *Frankliniella occidentalis* (Pergande) (Thysanoptera: Thripidae) from greenhouses in the Niagara region of Ontario, 907.
- Broatch, J.S., and R.S. Vernon.** Comparison of water pan traps and sticky traps for monitoring *Delia* spp. (Diptera: Anthomyiidae) in canola, 979.
- Brodeur, J., A. Bouchard, and G. Turcotte.** Potential of four species of predatory mites as biological control agents of the tomato russet mite, *Aculops lycopersici* (Massee) (Eriophyiidae), 1.
- Brooks, R.J.** See Bonduriensky and Brooks, 827.
- Brooks, S.E., and J.D. Shorthouse.** Biology of the rose stem Galler *Diptolepis nodulosa* (Hymenoptera: Cynipidae) and its associated component community in central Ontario, 1121.
- Burgin, S.G., and F.F. Hunter.** Evidence of honeydew feeding in black flies (Diptera: Simuliidae), 859.
- Cabana, J.** See Trudel et al., 197.
- Candau, J.-N.** See Jarry et al., 7.
- Cantin-Plante, N.** See Plowright et Catin-Plante, 915.
- Casagrande, R.A.** See Van Driesche et al., 601.
- Chew, P.S.** See Rhainds et al., 927.
- Childs, R.** See Van Driesche et al., 601.
- Coderre, D.** See Lalongé et al., 371.
- Côté, S.** See Huber et al., 959.
- DeClerck-Floate, R.** See Peschken et al., 949.
- Del Pilar Iglesias Calvin, M.** See Pintureau et al., 35.
- Denke, P.M.** See Blodgett et al., 377.

- Diehl, J.K., N.J. Holliday, C.J. Lindgren, and R.E. Roughley.** Insects associated with purple loosestrife, *Lythrum salicaria* L., in southern Manitoba, 937.
- Digweed, S.C., J.R. Spence, and D.W. Langor.** Exotic birch-leafmining sawflies (Hymenoptera: Tenthredinidae) in Alberta: Distributions, seasonal activities, and the potential for competition, 319.
- Dirzo, R.** See Morón-Ríos et al., 71.
- Duke, G.M.** See Goettel et al., 1059.
- Duncan, R.W.** See Mitchell et al., 745.
- Dwyer, E., D.J. Larson, and I.D. Thompson.** Oribatida (Acari) in balsam fir [*Abies balsamea* (L.)] forests of western Newfoundland, 151.
- Ebling, P.M., and W.J. Kaupp.** Pathogenicity of a nuclear polyhedrosis virus to forest tent caterpillar, *Malacosoma disstria* (Hübner) (Lepidoptera: Lasiocampidae), 195.
- Edwards, M.A., and W.D. Seabrook.** Evidence for an airborne sex pheromone in the Colorado potato beetle, *Leptinotarsa decemlineata*, 667.
- Eidt, D.C., and J.G. Stewart.** Effect on growth and root nodulation of clovers, *Trifolium* spp., by *Galleria mellonella* (L.) (Lepidoptera: Pyralidae) infected with *Steinernema carpocapsae* (Weiser) (Rhabdita: Steinernematidae) and its symbiont, *Xenorhabdus nematophilus* Poinar and Thomas, 205.
- Fields, P.G., and N.D.G. White.** Survival and multiplication of stored-product beetles at simulated and actual winter temperatures, 887.
- Fitzpatrick, S.M.** See Li and Fitzpatrick, 363.
- Galante, E.** See Verdú et al., 657.
- García, E.R., and M.E.E. Zamora.** New records of plant species used by adult Monarch butterflies *Danaus plexippus* L. (Lepidoptera: Nymphalidae: Danainae) during migration in Mexico, 375.
- Gardiner, M.G.T.** See Judd and Gardiner, 815.
- Gaul, S.O.** See Neil et al., 1079.
- Gayubo, S.F.** See Torres et al., 1067.
- Gerber, G.H.** Oviposition preferences of *Lygus lineolaris* (Palisot de Beauvois) (Heteroptera: Miridae) on four *Brassica* and two *Sinapis* species (Brassicaceae) in field cages, 855.
- Gerber, G.H.** See Leferink and Gerber, 777.
- Gillan, T.L., and D.H.S. Richardson.** The chalcid seed wasp, *Megastigmus nigrovariegatus* Hymenoptera: Torymidae, on *Rosa rugosa* Thunb. in Nova Scotia, 809.
- Gillespie, D.R.** See Opit et al., 221.
- Gnaspi, P.** See Peck and Gnaspi, 93.
- See Peck and Gnaspi, 769.
- Goerzen, D.W.** See Goettel et al., 1059.
- Goettel, M.S.** See Inglis et al., 171.
- Goettel, M.S., G.M. Duke, and D.W. Goerzen.** Pathogenicity of *Ascospaera larvis* to larvae of the alfalfa leafcutting bee, *Megachile rotundata*, 1059.
- Good, W.R.** See Story et al., 373.
- Grant, G.G.** See Silk et al., 1001.
- Grenier, S.** See Pintureau et al., 35.
- Gries, G.** See Rhainds et al., 199.
- See Rhainds et al., 927.
- Gries, G., R. Gries, and P.W. Schaefer.** Pheromone blend attracts nun moth, *Lymantria monacha* (Lepidoptera: Lymantriidae), in Japan, 1177.
- Guertin, C.** See Trudel et al., 197.
- Haack, R.A.** See Lawrence et al., 291.
- Hammond, H.E.J.** Arthropod biodiversity from *Populus* coarse woody material in north-central Alberta: A review of taxa and collection methods, 1009.
- Harvey, G.T.** Interspecific crosses and fertile hybrids among the coniferophagous *Choristoneura* (Lepidoptera: Tortricidae), 519.
- Headrick, D.H., T.S. Bellows, Jr., and T.M. Perring.** Behavior of female *Eretmocerus* sp. nr. *californicus* Howard (Aphelinidae: Hymenoptera) attacking *Bemisia argentifolii* Bellows and Perring (Aleyrodidae: Homoptera) on two native Californian weeds, 335.
- Helson, B.** See Nicol et al., 991.
- Helson, B.V.** See Wanner et al., 645.

- Holliday, N.J.** See Diehl et al., 937.
- Horton, D.R.** See Weissling et al., 637.
- Huber, J.T., S. Côté, and G. Boivin.** Description of three new *Anaphes* species (Hymenoptera: Mymaridae), egg parasitoids of the carrot weevil, *Listronotus oregonensis* (LeConte) (Coleoptera: Curculionidae), and redescription of *Anaphes sordidatus* Girault, 959.
- Hui, Y., and A. Bakke.** Development and reproduction of *Thanasisus formicarius* (L.) (Coleoptera: Cleridae) at three constant temperatures, 579.
- Hunter, F.F.** See Burgin and Hunter, 859.
- Inglis, G.D., D.L. Johnson, and M.S. Goettel.** Field and laboratory evaluation of two conidial batches of *Beauveria bassiana* (Balsamo) Vuillemin against grasshoppers, 171.
- Ivie, M.A.** See Blodgett et al., 377.
- Jaramillo, V.J.** See Morón-Ríos et al., 71.
- Jarry, M., J.-N. Candau, A. Roques, and B. Ycart.** Impact of emigrating seed chalcid, *Megastigmus spermatrophus* Wachtl (Hymenoptera: Torymidae), on seed production in a Douglas-fir seed orchard in France and modelling of orchard invasion, 7.
- Jobin, L.** See Retnakaran et al., 871.
- Johnson, D.L.** See Inglis et al., 171.
- Judd, G.J.R., and M.G.T. Gardiner.** Forecasting phenology of *Orthosia hibisci* Guenée (Lepidoptera: Noctuidae) in British Columbia using sex-attractant traps and degree-day models, 815.
- Kacem, N.** See Nénon et al., 789.
- Kaupp, W.J.** See Ebling and Kaupp, 195.
- Kays, S.J.** See Starr et al., 61.
- Kenis, M.** See West and Kenis, 831.
- Kennedy, G.G.** See Stewart et al., 21.
- Kortello, A.** See Pritchard and Kortello, 733.
- Kostyk, B.C.** See Wanner et al., 645.
- Kuenen, L.P.S.** See Silk et al., 1001.
- Lalongé, S., J.H. Redner, and D. Coderre.** First Canadian records of *Trochosa ruricola* (DeGeer), *Ostearius melanopygius* (O. Pickard-Cambridge), and *Dictyna decapirini* Kaston (Araneae: Lycosidae, Linyphiidae, Dictynidae, respectively), 371.
- Lamb, R.J.** See Palaniswamy et al., 43.
- Lamb, R.J., and P.A. Mackay.** Photoperiodism and life cycle plasticity of an aphid, *Macrosiphum euphorbiae* (Thomas), from central North America, 1035.
- Lamb, R.J., I.L. Wise, and P.A. Mackay.** Photoperiodism and seasonal abundance of an aphid, *Macrosiphum euphorbiae* (Thomas), in oilseed flax, 1049.
- Langor, D.W.** See Digweed et al., 319.
- Laplante, S.** See Bousquet and Laplante, 699.
- Larson, D.J.** Revision of North American *Agabus* Leach (Coleoptera: Dytiscidae): The *seriatus*-group, 105.
See Dwyer et al., 151.
See Puvanendran et al., 505.
- Lawrence, R.K., W.J. Mattson, and R.A. Haack.** White spruce and the spruce budworm: Defining the phenological window of susceptibility, 291.
- Leferink, J.H.M., and G.H. Gerber.** Development of adult and nymphal populations of *Lygus lineolaris* (Palisot de Beauvois), *L. elisus* Van Duzee, and *L. borealis* (Kelton) (Heteroptera: Miridae) in relation to seeding date and stage of plant development on canola (Brassicaceae) in southern Manitoba, 777.
- Le Lannic, J.** See Nénon et al., 789.
- Lenssen, A.W.** See Blodgett et al., 377.
- Lewis, T.M.** See Weissling et al. 637.
- Li, S.Y., and S.M. Fitzpatrick.** Responses of larval *Choristoneura rosaceana* (Harris) (Lepidoptera: Tortricidae) to a feeding stimulant, 363.
- Lindgren, C.J.** See Diehl et al., 937.
- Liu, Y.-B., and B.E. Tabashnik.** Visual determination of sex of diamondback moth larvae, 585.
- Loneragan, G.C.** See Silk et al., 1001.
- Lumaret, J.P.** See Verdú et al., 657.

- Mackay, P.A.** See Lamb and Mackay, 1035.
See Lamb et al., 1049.
- Marshall, S.A.** See Skevington and Marshall, 387.
- Marshall, V.G.** See Mitchell et al., 745.
- Mason, P.G.** See Adler and Mason, 81.
- Mattson, W.J.** See Lawrence et al., 291.
- McClay, A.S.** See Peschken et al., 949.
- McCorquodale, D.B., and R.E. Owen.** Allozyme variation, relatedness among progeny in a nest, and sex ratio in the leafcutter bee, *Megachile rotundata* (Fabricius) (Hymenoptera: Megachilidae), 211.
- McDonough, L.M.** See Weissling et al., 637.
- McRae, K.B.** See Neil et al., 1079.
- Mickle, R.** See West et al., 613.
- Mier Durante, M.P.** See Ortego et al., 1093.
- Mitchell, A.K., R.W. Duncan, T.A. Bown, and V.G. Marshall.** Origin and distribution of the yew big bud mite, *Cecidophyopsis psilaspis* (Nalepa), in British Columbia, 745.
- Morón-Ríos, A., V.J. Jaramillo, and R. Dirzo.** Species composition of root-feeding macroarthropods in a subalpine grassland associated with pine forest in Mexico, 71.
- Myles, T.G.** A second species of the drywood termite genus *Marginitermes* (Isoptera: Kalotermitidae), 757.
- Neil, K.A., S.O. Gaul, and K.B. McRae.** Control of the English grain aphid [*Sitobion avenae* (F.)] (Homoptera: Aphididae) and the oat-birdcherry aphid [*Rhopalosiphum padi* (L.)] (Homoptera: Aphididae) on winter cereals, 1079.
- Nénon, J.-P., N. Kacem, and J. Le Lannic.** Structure, sensory equipment, and secretions of the ovipositor in a giant species of Hymenoptera: *Megarhyssa atrata* F. (Ichneumonidae: Pimplinae), 789.
- Nicol, R.W., J.T. Arnason, B. Helson, and M.M. Abou-Zaid.** Effect of host and nonhost trees on the growth and development of the forest tent caterpillar, *Malacosoma disstria* (Lepidoptera: Lasiocampidae), 991.
- Nieto Nafria, J.M.** See Ortego et al., 1093.
- Norton, R.A.** See Perrot-Minnot and Norton, 691.
- O'Brien, C.W.** See Blodgett et al., 377.
- O'Keefe, S., T. Pike, and G. Poinar.** *Palaeolectochromus schaufussi* (gen.nov., sp.nov.), a new antlike stone beetle (Coleoptera: Scydmaenidae) from Canadian Cretaceous amber, 379.
- Onyabe, D.Y., and B.D. Roitberg.** The effect of conspecifics on oviposition site selection and oviposition behaviour in *Aedes togoi* (Theobald) (Diptera: Culicidae), 1173.
- Opit, G.P., B. Roitberg, and D.R. Gillespie.** The functional response and prey preference of *Feltiella acarissuga* (Vallot) (Diptera: Cecidomyiidae) for two of its prey: Male and female twospotted spider mites, *Tetranychus urticae* Koch (Acari: Tetranychidae), 221.
- Ortego, J., J.M. Nieto Nafria, and M.P. Mier Durante.** *Blanchardia poikila* gen.nov., sp.nov. (Hemiptera: Aphididae, Macrosiphini) in Mendoza, Argentina, 1093.
- Osborn, W.P.L.** See Boiteau and Osborn, 241.
- Owen, R.E.** See McCorquodale and Owen, 211.
- Packer, L., and J.S. Taylor.** FORUM: How many hidden species are there? An application of the phylogenetic species concept to genetic data for some comparatively well known bee "species", 587.
- Palaniswamy, P., R.J. Lamb, and R.P. Bodnaryk.** Antibiosis of preferred and non-preferred host-plants for the flea beetle, *Phyllotreta cruciferae* (Goeze) (Coleoptera: Chrysomelidae), 43.
- Palli, S.R.** See Retnakaran et al., 871.
- Pankiw, T.** Queen rearing by high and low queen mandibular pheromone responding worker honey bees (*Apis mellifera* L.), 679.
- Partridge, M.J., and J.H. Borden.** Evaluation of neem seed extract for control of the spruce aphid, *Elatobium abietinum* (Walker) (Homoptera: Aphididae), 899.
- Payne, N.** See Retnakaran et al., 871.
- Peck, S.B., and P. Gnaspini.** Review of the myrmecophilous *Ptomaphagus*, subgenus *Echinocoleus* (new status), of North America (Coleoptera: Leiodidae: Cholevinae: Ptomaphagini), 93.
- Peck, S.B., and P. Gnaspini.** *Ptomaphagus inyoensis* n.sp., a new microphthalmic montane beetle from California (Coleoptera: Leiodidae: Cholevinae: Ptomaphagini), 769.
- Pelletier, Y., et G. Bélanger.** Effet du rayonnement lumineux et de la vitesse de l'air ambiant sur la température corporelle des larves et des adultes du doryphore de la pomme de terre, 801.

- Perring, T.M.** See Headrick et al., 335.
- Perrot-Minnot, M.-J., and R.A. Norton.** Obligate thelytoky in oribatid mites: No evidence for *Wolbachia* inducement, 691.
- Peschken, D.P., R. DeClerck-Floate, and A.S. McClay.** *Cassida azurea* Fab. (Coleoptera: Chrysomelidae): Host specificity and establishment in Canada as a biological control agent against the weed *Silene vulgaris* (Moench) Garcke, 949.
- Pike, T.** See O'Keefe et al., 379.
- Pintureau, B., M. Del Pilar Iglesias Calvin, and S. Grenier.** Effectiveness of the second mating in a bisexual *Trichogramma* species and the first mating in a thelytokous *Trichogramma* species (Hymenoptera: Trichogrammatidae), 35.
- Plowright, C., et N. Cantin-Plante.** Le choix de la direction des trajets entre fleurs des bourdons (*Bombus*) butineurs, 915.
- Plowright, C.M.S., and R.C. Plowright.** The advantage of short tongues in bumble bees (*Bombus*)—Analyses of species distributions according to flower corolla depth, and of working speeds on white clover, 51.
- Plowright, C.M.S., R.C. Plowright, and P.H. Williams.** Replacement of *Bombus muscorum* by *Bombus pascuorum* in northern Britain?, 985.
- Plowright, R.C.** See Plowright and Plowright, 51.
See Plowright et al., 985.
- Poinar, G.** See O'Keefe et al., 379.
- Pree, D.J.** See Broadbent and Pree, 907.
- Primavera, M.J.** See Retnakaran et al., 871.
- Pritchard, G.** See Zloty and Pritchard, 251.
- Pritchard, G., and A. Kortello.** Roosting, perching, and habitat selection in *Argia vivida* Hagen and *Amphiargia abbreviatum* (Selys) (Odonata: Coenagrionidae), two damselflies inhabiting geothermal springs, 733.
- Puvanendran, D.C., D.J. Larson, and I.D. Thompson.** Collembola (Arthropoda) of balsam fir [*Abies balsamea* (L.)] forests of western Newfoundland, 505.
- Quednau, F.W.** Taxonomic notes on three black-bordered *Myzocallis* from oak in North America with descriptions of two new species (Homoptera: Aphididae), 1161.
- Redner, J.H.** See Lalongé et al., 371.
- Retnakaran, A.** See West et al., 613.
- Retnakaran, A., L.F.R. Smith, W.L. Tomkins, M.J. Primavera, S.R. Palli, N. Payne, and L. Jobin.** Effect of RH-5992, a nonsteroidal ecdysone agonist, on the spruce budworm, *Choristoneura fumiferana* (Lepidoptera: Tortricidae): Laboratory, greenhouse, and ground spray trials, 871.
- Rhainds, M., G. Gries, and P.S. Chew.** Adaptive significance of density-dependent ballooning by bagworm larvae, *Metisa plana* (Walker) (Lepidoptera: Psychidae), 927.
- Rhainds, M., G. Gries, and A. Saleh.** Sex-specific habitat utilization by bagworms (Lepidoptera: Psychidae), 199.
- Richardson, D.H.S.** See Gillan and Richardson, 809.
- Roitberg, B.** See Opit et al., 221.
- Roques, A.** See Jarry et al., 7.
- Roughley, R.E.** See Diehl et al., 937.
- Saeed, M.** See AliNiazee et al., 27.
- Saleh, A.** See Rhainds et al., 199.
- Sandoval, C.P.** See Vickery and Sandoval, 933.
- Seabrook, W.D.** See Edwards and Seabrook, 667.
- Severson, R.F.** See Starr et al., 61.
- Shorthouse, J.D.** See Brooks and Shorthouse, 1121.
- Silk, P.J., L.P.S. Kuenen, A.R. Alford, G.C. Lonergan, and G.G. Grant.** Sex pheromone components of the oak leaf shredder, *Croesia semipurpurana* (Kearfott) (Lepidoptera: Tortricidae), 1001.
- Skewington, J., and S.A. Marshall.** First record of a big-headed fly, *Eudorylas alternatus* (Cresson) (Diptera: Pipunculidae), reared from the subfamily Cicadellinae (Homoptera: Cicadellidae), with an overview of pipunculid-host associations in the Nearctic region, 387.
- Smith, L.F.R.** See Retnakaran et al., 871.

- Songa, J.M., and N.J. Holliday. Laboratory studies of predation of grasshopper eggs, *Melanoplus bivittatus* (Say), by adults of two species of *Pterostichus* Bonelli (Coleoptera: Carabidae), 1151.
- Spence, J.R. See Digweed et al., 319.
- Starr, C.K., D.D. Wilson, R.F. Severson, and S.J. Kays. Sexual dimorphism in the sweet potato weevil, *Cylas formicarius* (F.) (Coleoptera: Brentidae), 61.
- Stevenson, A.B., and E.S. Barszcz. A comparison of three methods of monitoring the carrot weevil, *Listronotus oregonensis* (LeConte) (Coleoptera: Curculionidae), 187.
- Stewart, H.C., and S.J. Walde. The dynamics of *Aphis pomi* De Geer (Homoptera: Aphididae) and its predator, *Aphidoletes aphidimyza* (Rondani) (Diptera: Cecidomyiidae), on apple in Nova Scotia, 627.
- Stewart, J.G. See Eidt and Stewart, 205.
- Stewart, J.G., G.G. Kennedy, and A.V. Sturz. Incidence of insecticide resistance in populations of Colorado potato beetle, *Leptinotarsa decemlineata* (Say) (Coleoptera: Chrysomelidae), on Prince Edward Island, 21.
- Story, J.M., W.R. Good, and L.J. White. First report of the establishment of *Cyphocleonus achates* (Fahraeus) (Coleoptera: Curculionidae) on spotted knapweed, *Centaurea maculosa* Lamarck, in the United States, 373.
- Sturz, A.V. See Stewart et al., 21.
- Sundaram, A. See West et al., 613.
- Sundaram, K.M.S. See West et al., 613.
- Tabashnik, B.E. See Liu and Tabashnik, 585.
- Taylor, J.S. See Packer and Taylor, 587.
- Tewksbury, L. See Van Driesche et al., 601.
- Thompson, D. See West et al., 613.
- Thompson, I.D. See Dwyer et al., 151.
See Puvanendran et al., 505.
- Tomkins, W.L. See Retnakaran et al., 871.
- Tormos, J. See Torres et al., 1067.
- Torres, F., S.F. Gayubo, J. Tormos, and J.D. Asis. Description of the mature larvae of three *Hoplitis* (Hymenoptera: Apoidea: Megachilidae), 1067.
- Trudel, R., É. Baucé, J. Cabana, and C. Guertin. Vulnerability of the fir coneworm, *Dioryctria abietivorella* (Grote) (Lepidoptera: Pyralidae), in different larval stages to the HD-1 strain of *Bacillus thuringiensis*, 197.
- Turcotte, G. See Brodeur et al., 1.
- Van Driesche, R.G., R. Childs, R.A. Casagrande, and L. Tewksbury. Establishment, distribution, and impact in southern New England of *Lathrolestes nigricollis* (Thompson) (Hymenoptera: Ichneumonidae), an introduced parasitoid of the birch leafminer, *Fenusa pusilla* (Lepeletier) (Hymenoptera: Tenthredinidae), 601.
- Verdú, J.R., J.P. Lumaret, and E. Galante. Biology of *Aphodius hyxos* Petrovitz (Coleoptera: Scarabaeoidea: Aphodiidae) and description of the third larval stage, 657.
- Vernon, R.S. See Broatch and Vernon, 979.
- Vickery, V.R. *Rubrogryllus coloratus*, a new cricket genus and species (Grylloptera: Gryllidae: Gryllinae) from Baja California, Mexico, 673.
- Vickery, V.R. Two new species in the genus *Diapheromera* (Phasmatoptera: Necroscoioidea: Heteronemiidae: Heteronemiinae) from Baja California, Mexico, 1141.
- Vickery, V.R., and C.P. Sandoval. *Timema bartmani* (Phasmatoptera: Timematodea: Timematidae), a new species from southern California, 933.
- Vrain, T.C. See West and Vrain, 229.
- Walde, S.J. See Stewart and Walde, 627.
- Wanner, K.W., B.V. Helson, and B.C. Kostyk. Foliar and systemic applications of neem seed extract for control of spruce budworm, *Choristoneura fumiferana* (Clem.) (Lepidoptera: Tortricidae), infesting black and white spruce seed orchards, 645.
- Webster, R.P., and C.-M. Yin. Effects of photoperiod and temperature on calling behaviour of the gypsy moth, *Lymantria dispar* L. (Lepidoptera: Lymantriidae), 843.
- Weissling, T.J., T.M. Lewis, L.M. McDonough, and D.R. Horton. Reduction in pear psylla (Homoptera: Psyllidae) oviposition and feeding by foliar application of various materials, 637.
- Weseloh, R.M. Orientation of *Calosoma sycophanta* L. (Coleoptera: Carabidae) in forests: Insights from visual responses to objects, 347.

- Weseloh, R.M.** Evidence for limited dispersal of larval gypsy moth, *Lymantria dispar* L. (Lepidoptera: Lymantriidae), 355.
- West, R.J., and M. Kenis.** Screening four exotic parasitoids as potential controls for the eastern hemlock looper, *Lambdina fiscellaria fiscellaria* (Guenée) (Lepidoptera: Geometridae), 831.
- West, R.J., D. Thompson, K.M.S. Sundaram, A. Sundaram, A. Retnakaran, and R. Mickle.** Efficacy of aerial applications of *Bacillus thuringiensis* Berliner and tebufenozide against the eastern hemlock looper (Lepidoptera: Geometridae), 613.
- West, R.J., and T.C. Vrain.** Nematode control of black army cutworm (Lepidoptera: Noctuidae) under laboratory and field conditions, 229.
- White, L.J.** See Story et al., 373.
- White, N.D.G.** See Fields and White, 887.
- Williams, P.H.** See Plowright et al., 985.
- Wilson, D.D.** See Starr et al., 61.
- Winchester, N.N.** FORUM: The arboreal superhighway: arthropods and landscape dynamics, 595.
- Wise, I.L.** See Lamb et al., 1049.
- Ycart, B.** See Jarry et al., 7.
- Yin, C.-M.** See Webster and Yin, 843.
- Zamora, M.E.E.** See García and Zamora, 375.
- Zloty, J., and G. Pritchard.** Larvae and adults of *Ameletus* mayflies (Ephemeroptera: Ameletidae) from Alberta, 251.

ERRATA

- Bousquet, Y., and S. Laplante.** Taxonomic review of the New World Pogonini (Coleoptera: Carabidae), 1179.
- LeSage, L., and P. Paquin.** Identification keys for *Aphthona* flea beetles (Coleoptera: Chrysomelidae) introduced in Canada for the control of spurge (*Euphorbia* spp., Euphorbiaceae), 201.

